

# Comparing LANDFIRE fuel representation systems & their application in estimating fire effects

Josh Hyde<sup>1</sup>, Eva Strand<sup>2</sup>, Andrew Hudak<sup>3</sup>

<sup>1</sup>Smoke Program Coordinator, University of Idaho College of Natural Resources Department of Forest, Rangeland, and Fire Sciences, Moscow, ID, United States


<sup>2</sup>Assistant professor, University of Idaho College of Natural Resources Department of Forest, Rangeland, and Fire Sciences, Moscow, ID, United States

<sup>3</sup> Research Forester, Rocky Mountain Research Station, United States Forest Service, Moscow, ID, United States

**Abstract:** Managers and researchers are often tasked with estimating the impacts of wildland fire on landscapes. One data source for performing these assessments is the fire effects fuel layers available from LANDFIRE; the Fuels Classification Characterization System (FCCS) and the Fuel Loading Model (FLM) spatial layers. The two spatial layers were developed independently of one another. This study evaluates the differences between LANDFIRE FCCS (FCCS<sub>L</sub>) and LANDFIRE FLM (FLM<sub>L</sub>) layers with regards to fuel loading, and the subsequent differences in consumption and fire effects when modeled with the Wildland Fire Assessment Tool (WFAT). A case study in mixed conifer northern Idaho forest is presented. Results indicated estimated duff loading was likely to be higher in the LANDFIRE FCCS fuel layer, estimated 1000 hr fuel loading lower in the FLM layer. Shrub loadings were greater in the FLM layer compared to the FCCS layer.


## Methods

To evaluate the potential differences in WFAT outputs given FCCS<sub>L</sub> and FLM<sub>L</sub> inputs the authors examined a 700-ha study area centered on Moscow Mountain in Latah County Idaho. The area is dominated by mixed conifer forest tree species including ponderosa pine, Douglas-fir occurring on more xeric southern and western slopes and grand fir and cedar/hemlock habitat types occur on the more mesic northern and eastern aspects (Cooper et al. 1991).




To compare fuel loading, consumption, and fire effects FCCS<sub>L</sub> and FLM<sub>L</sub> data was compared with measured fuel loadings and estimated shrub loadings from 87 randomly-placed field sampling plots collected in 2009 (Hudak et al. 2012) (*above*). Loading data including woody fuels, litter, duff, herb, and estimated shrub loading using Brown's (1981) methodology.

LANDFIRE Refresh 2008 FCCS<sub>L</sub> and FLM<sub>L</sub> layers were downloaded for the area. The Wildland Fire Assessment Tool (WFAT) (Hamilton et al. 2012) was then used to simulate fuel consumption, emissions, and soil heating. WFAT is a spatial analysis tool employing Flamap (Finney 2006, Nelson 2000, Rothermel 1972, Van Wagner 1977) algorithms for fire behavior, and First Order Fire Effects Model (FOFEM) (Abini and Reinhardt 1995; Albini et al. 1995; Albini and Reinhardt 1997, Reinhardt 2003) algorithms for fire effects. Fuel loading, Consumption, and emissions estimates were compared for the measured fuel data, FCCS<sub>L</sub> fuel data, and FLM<sub>L</sub> fuel data using analysis of variance (R Core Team 2012). Parameters to populate WFAT (*below*) were taken from LANDFIRE.



## Results & Conclusions

- FCCS<sub>L</sub> duff loading was greater than either FLM<sub>L</sub> or measured data (*below*).
- For litter, 10, and 100-hour fine woody fuels, the measured loading was greater than FCCS<sub>L</sub> and FLM<sub>L</sub> layers.
- One-hour fine woody fuel loading in the FCCS<sub>L</sub> was higher than that of FLM<sub>L</sub>.
- Thousand-hour fuel loading was less for LANDFIRE FLM<sub>L</sub> than FCCS<sub>L</sub> or measured data.
- Herbaceous and shrub loadings were greater for FLM<sub>L</sub> than measured herb loadings or estimated shrub loadings. The remaining comparisons were not significantly different from each other.
- The higher duff loadings in FCCS<sub>L</sub> resulted in more biomass consumed, and subsequently greater emissions and surface temperatures.
- Lower duff and litter loading in the FLM<sub>L</sub> resulted in greater mineral soil exposure.
- Fuel loading values for the study site and LANDFIRE layers were within ranges observed by other researcher for of North-western Rocky Mountain Fuels with the exception of duff. Duff loadings tended to be lower for our research site and the FLM<sub>L</sub> than other observed studies while FCCS<sub>L</sub> duff loading was closer to observed values (Hille & Stephens 2005, Youngblood et al. 2008, Reinhardt et al. 1991).
- LANDFIRE FCCS and FLM layers provide fuel information where little or no existing information may be available. However, a customized fuel layer, if data is available, is likely to represent the landscape more accurately than an unaltered FCCS<sub>L</sub> or FLM<sub>L</sub> layer.



## References:

- Albini FA, Reinhardt ED (1995) Modeling ignition and burning rate of large woody natural fuels. *International Journal of Wildland Fire*, 5, 81-91.
- Albini FA, Brown JK, Reinhardt ED, Ottmar RD (1995) Calibration of a large fuel burnout model. *International Journal of Wildland Fire*, 5, 173-192.
- Albini, FA, Reinhardt ED (1997) Improved Calibration of a Large Fuel Burnout Model. *International Journal of Wildland Fire*, 7, 21-28.
- Brown JK (1981) Bulk densities of nonuniform surface fuels and their application to fire modeling. *Forest Science*, 27, 677-683.
- Cooper SV, Neilman KE, & Roberts DW (1991) Forest habitat types of northern Idaho: A second approximation. Gen. Tech. Rep. INT-236. Ogden, UT, U.S. Department of Agriculture, Forest Service, Intermountain Forest & Range Exp. Stn. 143 p.
- Finney MA (2006) An overview of FlamMap modeling capabilities. In: Andrews PL, Butler BW, comps. *Fuels Management—How to Measure Success: Conference Proceedings*. 2006 March 28-30; Portland, OR. Proc. RMRS-P-41. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 809 pp.
- Hamilton D, Jones J, Hann W (2012) Wildland Fire Assessment Tool (WFAT) for ArcGIS 10 (version 2.2.0). National Interagency Fuels Technology Transfer. [online] accessed January 7 2013. Available at <www.nifft.gov>.
- Hille MG, Stephens SI (2005) Mixed Conifer Forest Duff Consumption during Prescribed Fires: Tree Crown Impacts. *Forest Science*, 51, 417-424.
- Hudak AT, Strand EV, Vierling LA, Byrne JC, Eitel JH, Martinuzzi S, and Falkowski AU (2012) Quantifying aboveground forest carbon pools and fluxes from repeat LiDAR surveys. *Remote Sensing of Environment*, 123, 25-40.
- Lutes DC, Keane RE, and Caratti JF (2009) A surface fuels classification for estimating fire effects. *International Journal of Wildland Fire*, 18, 802-814.
- Nelson RM (2000) Prediction of diurnal change in 10-h fuel stick moisture content. *Canadian Journal of Forest Research*, 30, 1071-1087.
- Ottmar RD, Sandberg DV, Riccardi CL, and Pritchard SJ (2007) An overview of the Fuel Characteristic Classification System - Quantifying, classifying, and creating fuelbeds for resource planning. *Canadian J. of Forest Research*, 3, 2389-2393.
- R Core Team (2012) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- Reinhardt ED (2003) Using FOFEM 5.0 to estimate tree mortality, fuel consumption, smoke production and soil heating from wildland fire. In: *Proceedings of the Second International Wildland Fire Ecology and Fire Management Congress and Fifth Symposium on Fire and Forest Meteorology*, November 16-20, 2003, Orlando, FL. American Meteorological Society. P5-2.
- Reinhardt ED, Brown JK, Fischer WC, Graham RT (1991) Woody Fuel and Duff Consumption by Prescribed Fire in Northern Idaho Mixed Conifer Logging Slash. Research Paper INT-443. USDA Forest Service Intermountain Res. Stn 26 pgs.
- Rothermel RC (1972) A mathematical model for predicting fire spread in wildland fuels. General Technical Report INT-115. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest & Range Exp. Stn. 50 p.
- Van Wagner CE (1977) Conditions for the start and spread of crownfire. *Canadian Journal of Forest Research*, 7, 23-24.
- Youngblood A, Wright CS, Ottmar RD, McIver JD (2008) Changes in fuelbed characteristics and resulting fire potentials after fuel reduction treatments in dry forests of the Blue Mountains North Eastern Oregon. *For. Eco. & Mgt.* 255, 3151-3169.